

# Package ‘BeyondBenford’

January 19, 2019

**Type** Package

**Title** Compare the Goodness of Fit of Benford's and Blondeau Da Silva's Digit Distributions to a Given Dataset

**Version** 1.1

**Date** 2019-01-19

**Author** Blondeau Da Silva Stephane

**Maintainer** Blondeau Da Silva Stephane <Stephane.Blondeau-Da-Silva@ac-limoges.fr>

**Description** Allows to compare the goodness of fit of Benford's and Blondeau Da Silva's digit distributions in a dataset. It is used to check whether the data distribution is consistent with theoretical distributions highlighted by Blondeau Da Silva or not (through the `dat.distr()` function): this ideal theoretical distribution must be at least approximately followed by the data for the use of Blondeau Da Silva's model to be well-founded. It also enables to plot histograms of digit distributions, both observed in the dataset and given by the two theoretical approaches (with the `digit.ditr()` function). Finally, it proposes to quantify the goodness of fit via Pearson's chi-squared test (with the `chi2()` function).

**License** GPL-2

**NeedsCompilation** no

**Repository** CRAN

**Date/Publication** 2019-01-19 14:40:03 UTC

## R topics documented:

|                        |    |
|------------------------|----|
| BeyondBenford-package  | 2  |
| address_AixesurVienne  | 3  |
| address_Limoges        | 4  |
| address_PierreBuffiere | 4  |
| Benf.val               | 5  |
| Blon.val               | 6  |
| census                 | 7  |
| chi2                   | 7  |
| dat.distr              | 8  |
| digit.distr            | 10 |
| obs.numb.dig           | 12 |
| theor.distr.val        | 13 |

---

BeyondBenford-package *Compare the goodness of fit of Benford's and Blondeau Da Silva's digit distributions to a given dataset*

---

### Description

The purpose of this package is to compare the goodness of fit of Benford's and Blondeau Da Silva's digit distributions in a dataset. The package is used to check whether the data distribution is consistent with theoretical distributions highlighted by Blondeau Da Silva or not (through the function 'dat.distr'): this ideal theoretical distribution must be at least approximately followed by the data for the use of Blondeau Da Silva's model to be well-founded. It also enables to plot histograms of digit distributions, both observed in the dataset and given by the two theoretical approaches (with the function 'digit.ditr'). Finally, it proposes to quantify the goodness of fit via Pearson's chi-squared test (with the function 'chi2').

### Author(s)

Blondeau Da Silva

Maintainer: Blondeau Da Silva

### References

- F. Benford (1938). The law of anomalous numbers. *Proceedings of the American Philosophical Society*, 78:127-131.
- A. Berger and T. Hill (2015). *An introduction to Benford's Law*. Princeton University Press, Princeton, NJ. ISSN/ISBN: 978-0-691-16306-2.
- S. Blondeau Da Silva (2019). Benford or not Benford: a systematic but not always well-founded use of an elegant law in experimental fields. *Communications in Mathematics and Statistics*. In press.
- S. Blondeau Da Silva (2018). Benford or not Benford: new results on digits beyond the first. <https://arxiv.org/abs/1805.01291>.
- T. Hill (1995). The significant-digit phenomenon. *The American Mathematical Monthly*, 102(4):322-327.
- S. J. Miller, editor (2015). *Benford's Law: Theory and Applications*. Princeton University Press, Princeton, NJ. ISSN/ISBN: 978-0-691-14761-1.
- R. Newcomb (1881). Note on the frequency of use of the different digits in natural numbers. *American Journal of Mathematics*, 4:39-40.
- K. Pearson (1900). On the criterion that a given system of deviations from the probable in the case of a correlated system of variables is such that it can be reasonably supposed to have arisen from random sampling. *Philosophical Magazine*, 50(302):157-175.

**Examples**

```
data(address_PierreBuffiere)
data(census)
data(address_AixesurVienne)

dat.distr(address_PierreBuffiere,nchi=6)
dat.distr(census,theor=0,nclass=100,dig=3)
dat.distr(address_AixesurVienne,upbound=75)

digit.distr(address_AixesurVienne,mod="ben&blo",upbound=75)
digit.distr(address_PierreBuffiere,mod="blo",dig=2)

chi2(address_PierreBuffiere,dig=2,pval=1)
chi2(address_PierreBuffiere,dig=2,pval=1,mod="blo")
```

---

address\_AixesurVienne *Street addresses of Aix-sur-Vienne*

---

**Description**

Street addresses of Aix-sur-Vienne, a town of approximately 5800 inhabitants in Haute-Vienne (France).

**Usage**

```
address_AixesurVienne
```

**Format**

A factor containing all 1911 existing street address numbers.

**Source**

From an open platform for French public data:

<https://www.data.gouv.fr/fr/datasets/base-d-adresses-nationale-ouverte-bano/> (<http://bano.openstreetmap.fr/data/>).

---

address\_Limoges      *Street addresses of Limoges*

---

**Description**

Street addresses of Limoges, a city of approximately 133600 inhabitants in Haute-Vienne (France).

**Usage**

address\_Limoges

**Format**

A factor containing all 35975 existing street address numbers.

**Source**

From an open platform for French public data:

<https://www.data.gouv.fr/fr/datasets/base-d-adresses-nationale-ouverte-bano/> (<http://bano.openstreetmap.fr/data/>).

---

address\_PierreBuffiere  
*Street addresses of Pierre-Buffiere*

---

**Description**

Street addresses of Pierre-Buffiere, a small town of approximately 1200 inhabitants in Haute-Vienne (France).

**Usage**

address\_PierreBuffiere

**Format**

A factor containing all 346 existing street address numbers.

**Source**

From an open platform for French public data:

<https://www.data.gouv.fr/fr/datasets/base-d-adresses-nationale-ouverte-bano/> (<http://bano.openstreetmap.fr/data/>).

---

|          |                         |
|----------|-------------------------|
| Benf.val | <i>Benford's values</i> |
|----------|-------------------------|

---

**Description**

The function returns Benford's probability that a figure is at a given position.

**Usage**

```
Benf.val(fig, dig = 1)
```

**Arguments**

|     |   |
|-----|---|
| fig | The considered figure.                            |
| dig | The chosen position of the digit (from the left). |

**Value**

The function returns Benford's probability.

**Author(s)**

Blondeau Da Silva St'e'phane

**References**

F. Benford (1938). The law of anomalous numbers. Proceedings of the American Philosophical Society, 78:127-131.

T. Hill (1995). The significant-digit phenomenon. The American Mathematical Monthly, 102(4):322-327.

R. Newcomb (1881). Note on the frequency of use of the different digits in natural numbers. American Journal of Mathematics, 4:39-40.

**Examples**

```
Benf.val(7, dig = 2)
```

---

`Blon.val`*Blondeau Da Silva's values*

---

**Description**

The function returns Blondeau Da Silva's probability that a figure is at a given position (once the associated upper bound has been specified).

**Usage**

```
Blon.val(upperbound, fig, dig = 1)
```

**Arguments**

|                         |  |
|-------------------------|--|
| <code>upperbound</code> | A positive integer, which characterizes the data. All (or most) of the data are lower than this "upper bound". |
| <code>fig</code>        | The considered figure.   |
| <code>dig</code>        | The chosen position of the digit (from the left).  |

**Value**

The function returns Blondeau Da Silva's probability.

**Author(s)**

Blondeau Da Silva Stephane

**References**

S. Blondeau Da Silva (2019). Benford or not Benford: a systematic but not always well-founded use of an elegant law in experimental fields. Communications in Mathematics and Statistics. In press.

S. Blondeau Da Silva (2018). Benford or not Benford: new results on digits beyond the first. <https://arxiv.org/abs/1805.01291>.

**Examples**

```
Blon.val(825, 5, dig = 3)
```

---

|        |                       |
|--------|-----------------------|
| census | <i>Alabama census</i> |
|--------|-----------------------|

---

**Description**

Populations in Alabama cities and towns.

**Usage**

```
census
```

**Format**

A data frame containing the populations of all 460 Alabama cities or towns (dimension: one row and 460 columns).

**Source**

From the United States Census Bureau:

<https://factfinder.census.gov/faces/tableservices/jsf/pages/productview.xhtml?src=bkmk>.

---

|      |                                   |
|------|-----------------------------------|
| chi2 | <i>Pearson's chi-squared test</i> |
|------|-----------------------------------|

---

**Description**

It is a test of goodness of fit to find out whether the distribution of first (second, third or fourth) digit in the studied data differs from two theoretical distributions (that of Benford and that of Blondeau Da Silva) or not. The null hypothesis states that the studied distribution is consistent with the considered theoretical distribution.

**Usage**

```
chi2(dat, mod = "ben", upbound = ceiling(max(dat)), dig = 1, pval = 0)
```

**Arguments**

|         |  |
|---------|--|
| dat     | The considered dataset, a data frame containing non-zero real numbers.   |
| mod     | If mod="ben", the theoretical distribution considered is that of Benford, else it is Blondeau Da Silva's ones which is chosen. |
| upbound | A positive integer, which characterizes the data. All (or most) of the data are lower than this "upper bound".                 |
| dig     | The chosen position of the digit (from the left).  |
| pval    | If pval=0, the p-value is not returned, else it is available.  |

**Value**

A data frame containing the Pearson chi-squared statistic (and the associated p-value if requested).

**Note**

This warning message can appear: NAs introduced during the automatic conversion. This is due to the fact that some data are not numerical in the entered dataset. Non numerical values and zeros are not counted.

**Author(s)**

Blondeau Da Silva Stephane

**References**

K. Pearson (1900). On the criterion that a given system of deviations from the probable in the case of a correlated system of variables is such that it can be reasonably supposed to have arisen from random sampling. *Philosophical Magazine*, 50(302):157-175.

**Examples**

```
data(address_PierreBuffiere)
chi2(address_PierreBuffiere,dig=2,pval=1)
chi2(address_PierreBuffiere,dig=2,pval=1,mod="blo")
```

---

dat.distr

*Data distribution*

---

**Description**

The function returns the histogram of the data. It can also plot one of the Blondeau Da Silva's theoretical distributions (thanks to an "upper bound"): this ideal theoretical distribution must be at least approximately followed by the data for the use of Blondeau Da Silva's model to be well-founded. A specific chi-squared statistic can also be computed to find out whether the data distribution is consistent with the theoretical distribution or not.

**Usage**

```
dat.distr(dat, xlab = "data", ylab = "Frequency", main = "Distribution of data",
  theor = TRUE, nclass = 50, col = "lightblue", conv = 0, upbound = ceiling(max(dat)),
  dig = 1, colt = "red", ylim = NULL, border = "blue", nchi = 0, legend = TRUE,
  bg.leg = "gray85")
```



**Arguments**

|         |   |
|---------|---|
| dat     | The considered dataset, a data frame containing non-zero real numbers.  |
| xlab    | The x-axis label.   |
| ylab    | The y-axis label.   |
| main    | The title of the graph.   |
| theor   | If theor=TRUE Blondeau Da Silva's theoretical distribution is plotted, otherwise only the histogram is represented.   |
| nclass  | A strictly positive integer: the number of classes in the histogram.  |
| col     | The color used to fill the bars of the histogram. NULL yields unfilled bars.  |
| conv    | If conv=1, all values of the dataset are multiplied by $10^k$ where k is the smallest positive integer such that all non-zero numerical values in the newly multiplied data frame have an absolute value greater than or equal to 1.  |
| upbound | A positive integer, which characterizes the data. All (or most) of the data are lower than this "upper bound".  |
| dig     | The chosen position of the digit (from the left).   |
| colt    | The color used to plot Blondeau Da Silva's theoretical distribution.  |
| ylim    | A two-components vector: the range of y values.   |
| border  | The color of the border around the bars.  |
| nchi    | A positive integer: the number of classes for values from $10^{(p-1)}$ to $\max(\max(\text{data}), \text{upbound})$ . If nchi>0, the function returns the chi-squared statistic (with nchi-1 degrees of freedom) of goodness of fit determined by the different classes. The null hypothesis states that the studied distribution is consistent with the considered theoretical distribution. |
| legend  | If legend=TRUE, the legend is displayed.  |
| bg.leg  | The background color for the legend box.  |

**Value**

The histogram of the data along with optional Blondeau Da Silva's theoretical distributions and a data frame containing the chi-squared statistic and its associated p-value if requested.

**Note**

This warning message can appear: NAs introduced during the automatic conversion. This is due to the fact that some data are not numerical in the entered dataset. Non numerical values and zeros are not counted.

**Author(s)**

Blondeau Da Silva Stephane

## References

S. Blondeau Da Silva (2019). Benford or not Benford: a systematic but not always well-founded use of an elegant law in experimental fields. *Communications in Mathematics and Statistics*. In press.

S. Blondeau Da Silva (2018). Benford or not Benford: new results on digits beyond the first. <https://arxiv.org/abs/1805.01291>.

K. Pearson (1900). On the criterion that a given system of deviations from the probable in the case of a correlated system of variables is such that it can be reasonably supposed to have arisen from random sampling. *Philosophical Magazine*, 50(302):157-175.

## Examples

```
data(address_PierreBuffiere)
dat.distr(address_PierreBuffiere,nchi=6)

data(census)
dat.distr(census,theor=0,nclass=100,dig=3)

data(address_AixesurVienne)
dat.distr(address_AixesurVienne,upbound=75)
```

---

digit.distr

*Distribution of figures in a given position*

---

## Description

The function returns histograms of distribution of figures in a given position: (i) in the dataset, (ii) due to Benford, (iii) due to Blondeau Da Silva.

## Usage

```
digit.distr(dat, mod = "ben", upbound = ceiling(max(dat)), dig = 1,
col = c("#FFFFAA", "#AAFFAA"), colbl = c("#FFFFAA", "#AAFFFF"),
colbebl = c("#FFFFAA", "#AAFFAA", "#AAFFFF"), main = "Distribution of digits",
legend = TRUE, leg = c("Observed", "Benford"),
legbebl = c("Observed", "Benford", "Blondeau"), legbl = c("Observed", "Blondeau"))
```

## Arguments

|         |  |
|---------|--|
| dat     | The considered dataset, a data frame containing non-zero real numbers.   |
| mod     | If mod="ben", the data histogram and that of Benford are displayed, if mod="ben&blo", the data histogram, that of Benford and that of Blondeau Da Silva are plotted, and otherwise the data histogram and that of Blondeau Da Silva are given. |
| upbound | A positive integer, which characterizes the data. All (or most) of the data are lower than this "upper bound".   |

|         |  |
|---------|--|
| dig     | The chosen position of the digit (from the left).  |
| col     | A vector containing two colors used to fill the bars of the histogram, if mod="ben".       |
| colbebl | A vector containing three colors used to fill the bars of the histogram, if mod="ben&blo". |
| colbl   | A vector containing two colors used to fill the bars of the histogram, if the latter case. |
| main    | The title of the graph.  |
| legend  | If legend=TRUE, the legend is displayed.   |
| leg     | A two-components vector containing text appearing in the legend, if mod="ben".             |
| legbebl | A three-components vector containing text appearing in the legend, if mod="ben&blo".       |
| legbl   | A two-components vector containing text appearing in the legend, if the latter case.       |

**Value**

Histograms of distribution of figures in a given position: (i) in the dataset, (ii) due to Benford, (iii) due to Blondeau Da Silva.

**Note**

This warning message can appear: NAs introduced during the automatic conversion. This is due to the fact that some data are not numerical in the entered dataset. Non numerical values and zeros are not counted.

**Author(s)**

Blondeau Da Silva Stephane

**References**

- F. Benford (1938). The law of anomalous numbers. *Proceedings of the American Philosophical Society*, 78:127-131.
- S. Blondeau Da Silva (2019). Benford or not Benford: a systematic but not always well-founded use of an elegant law in experimental fields. *Communications in Mathematics and Statistics*. In press.
- S. Blondeau Da Silva (2018). Benford or not Benford: new results on digits beyond the first. <https://arxiv.org/abs/1805.01291>.
- T. Hill (1995). The significant-digit phenomenon. *The American Mathematical Monthly*, 102(4):322-327.
- R. Newcomb (1881). Note on the frequency of use of the different digits in natural numbers. *American Journal of Mathematics*, 4:39-40.

**Examples**

```
data(address_AixesurVienne)
digit.distr(address_AixesurVienne,mod="ben&blo",upbound=75)
```

```
data(address_PierreBuffiere)
digit.distr(address_PierreBuffiere,mod="blo",dig=2)
```

---

|              |   |
|--------------|---|
| obs.numb.dig | <i>Frequency of each figure at a given position</i> |
|--------------|---|

---

**Description**

The function returns the frequencies of each figure at a given position in the considered dataset.

**Usage**

```
obs.numb.dig(dat, dig = 1)
```

**Arguments**

|     |  |
|-----|--|
| dat | The considered dataset, a data frame containing non-zero real numbers. |
| dig | The chosen position of the digit (from the left).                      |

**Value**

A vector containing the frequencies of each figure in ascending order. Its length is 9 if dig=1 (the figures ranging from 1 to 9) and 10 if dig>1 (the figures ranging from 0 to 9).

**Note**

This warning message can appear: NAs introduced during the automatic conversion. This is due to the fact that some data are not numerical in the entered dataset. Non numerical values and zeros are not counted.

**Author(s)**

Blondeau Da Silva Stephane

**Examples**

```
data(census)
obs.numb.dig(census, dig=2)
```

---

|                 |                                 |
|-----------------|---------------------------------|
| theor.distr.val | <i>Theoretical distribution</i> |
|-----------------|---------------------------------|

---

**Description**

The function returns the theoretical probability distribution described by Blondeau Da Silva for data. If the dataset follows this particular distribution well enough, it enables not to use Benford's values of first (second, third or fourth) digit distribution but rather Blondeau Da Silva's ones. The distribution depends on an upper bound, which characterizes the data.

**Usage**

```
theor.distr.val(upbound, dig = 1)
```

**Arguments**

|         |  |
|---------|--|
| upbound | A positive integer, which characterizes the data. All (or most) of the data are lower than this "upper bound". |
| dig     | The chosen position of the digit (from the left).  |

**Value**

The function returns a vector contening the probability distribution of the model determined by the upper bound value.

**Author(s)**

Blondeau Da Silva Stephane

**References**

S. Blondeau Da Silva (2019). Benford or not Benford: a systematic but not always well-founded use of an elegant law in experimental fields. Communications in Mathematics and Statistics. In press.

S. Blondeau Da Silva (2018). Benford or not Benford: new results on digits beyond the first. <https://arxiv.org/abs/1805.01291>.

**Examples**

```
theor.distr.val(10)
```

# Index

`address_AixesurVienne`, 3  
`address_Limoges`, 4  
`address_PierreBuffiere`, 4

`Benf.val`, 5  
`BeyondBenford` (`BeyondBenford-package`), 2  
`BeyondBenford-package`, 2  
`Blon.val`, 6

`census`, 7  
`chi2`, 7

`dat.distr`, 8  
`digit.distr`, 10

`obs.numb.dig`, 12

`theor.distr.val`, 13